

Two-colour photometry of the binary planetary nebula nuclei UU Sagitte and V477 Lyrae: oversized secondaries in post-common-envelope binaries

M. Afşar^{1*} and C. İbanoğlu¹

¹*Department of Astronomy and Space Sciences, Faculty of Science, Ege University, 35100 Bornova, İzmir, Turkey*

Accepted 2008 September 3. Received 2008 September 3; in original form 2008 April 23

ABSTRACT

We present new V - and R -passband CCD photometry of UU Sge and V477 Lyr, the eclipsing binary nuclei of the planetary nebulae Abell 63 and Abell 46, respectively. We have performed a simultaneous analysis of VR light-curves and estimated the effective temperatures for the primary and secondary stars to be $78\,000 \pm 3000$ and 6136 ± 240 K for UU Sge, $49\,500 \pm 4500$ and 3874 ± 350 K for V477 Lyr. We have also reanalysed the previously measured radial velocities and combined the results with those obtained from the analysis of the light curves to derive absolute parameters of the components. The secondary stars have larger radii than expected from their main-sequence counterparts at the same masses. We have determined the post-common envelope ages and the thermal time scales of the systems and examined the possible reasons of expanded radius of the secondary components, together with some selected post-common envelope binaries. We conclude that the secondary components of the nuclei of the planetary nebulae are still out of thermal equilibrium along with two post-common envelope systems: HS 1136+6646 and RE 1016-053. For other systems, magnetic activity has been suggested as the more plausible reason for their expanded radii. We have also estimated the common-envelope efficiency parameters of UU Sge and V477 Lyr.

Key words: binaries: close – binaries: eclipsing – stars: evolution – stars: individual:UU Sge – stars: individual:V477 Lyr – stars: AGB and post-AGB

1 INTRODUCTION

The classical approach to planetary nebulae (PNe) is that they are envelopes ejected by single stars ($0.8 \leq M/M_{\odot} \leq 10$) at the end of the asymptotic giant branch (AGB) stage. However, up to date, it has been reported that at least 10-15 per cent of planetary nebula nuclei (PNNi) are very close binaries with periods in the range of hours to a few days (Bond & Grauer 1987; Bond & Livio 1990; Bond 2000). Recently, some radial velocity surveys have been carried out in order to search for new variable PNNi, which have periods too long to be detected using photometry (e.g. De Marco et al. 2004; Afşar & Bond 2005). The existence of these objects has been explained by means of common envelope (CE) ejection by several authors (Paczynski 1976; Iben & Livio 1993; Iben & Tutukov 1993). According to CE evolution theories, the atmosphere of the primary component will expand during the red giant branch (RGB) or AGB evolution, and depending on the initial separation of the binary system, both components will be engulfed by extended atmosphere of the primary. The gravitational drag forces arise due to orbital motion of the secondary component around the primary's core (spiral-in process) will cause orbit to shrink dramatically. Because of the conservation of total angular momentum, the orbital angular momentum will be transferred to the CE and as a result the envelope is spun up and ejected from the system. Thus, the outcome will be a close binary as a nucleus of a planetary nebula. The efficiency of which orbital energy is deposited into CE ejection is defined as “efficiency parameter”, denoted as α_{CE} . Our knowledge on the CE evolution is highly limited due to the difficulties in determining of stellar and nebular parameters. However, the central stars of the PNe, which are also members of eclipsing binary systems, are the direct evidence of the existence of the CE phase and provide an opportunity to drive accurate absolute parameters of these objects.

There are a number of these objects and only four of them are eclipsing binaries: UU Sge, V477 Lyr, BE UMa and SuWt 2. The other close binary nuclei are known to be as binaries either due to reflection effect (see De Marco et al. 2008 for a detailed study) or ellipsoidal variation they show in their light curves, or they are just single-lined spectroscopic binaries.

The eclipsing binary nuclei UU Sge, the nucleus of Abell 63, and V477 Lyr, the nucleus of Abell 46, were previously studied by several authors. The eclipsing nature of UU Sge was first reported by Hoffleit in 1932. After three decades the presence of a low-surface brightness nebula was discovered by Abell (1966), and Bond (1976) showed the connection

between PN and UU Sge. The first photoelectric photometric observations of the system were made by Bond et al. (1978). They obtained a light curve in B -passband and indicated that the system has a total eclipse at the primary conjunction. They also noted the extreme heating effect in the system due to large temperature difference between the components. From the solution of the light curve they concluded that the system contains an sdO primary of $\sim 0.9 M_{\odot}$ with a temperature of about 35 000 K and a dK secondary of $\sim 0.7 M_{\odot}$. They also emphasized that the observations of the system was contaminated by the nearby optical companion $\sim 4''$ (later to be measured as $\sim 2''.8$ by Walton et al. 1993) to the east of UU Sge and they could not determine the true depth of primary minimum accurately. However, they estimated the contamination of the optical companion from the primary minimum observations and suggested that the actual depth of the primary eclipse should be about 4.3 mag.

Later, Budding & Kopal (1980) obtained the V -passband light curve of the system. They found the O-type primary component has a temperature of not much higher than 30 000 K and the secondary component is around 6000 K. Van Buren et al. (1980) discovered the soft X-ray emission feature of UU Sge and suggested that the observed X-ray originates in the corona of the late-type star.

Walton et al. (1993) made spectroscopic observations of the system at several phases. Using the spectrum taken during the primary minimum they estimated the spectral type of the secondary component as G7 V. From the spectra obtained at out-of-eclipse they deduced that the exciting hot central star is an sdO type PN nucleus with a temperature of around 50 000 K.

The first V - and I -passband CCD photometry was made by Pollacco & Bell (1993, hereafter PB93). However, in I -passband, they only obtained the primary minimum since they aimed to use the primary eclipse to determine an accurate temperature for the secondary. Using the medium-resolution spectroscopy, they obtained the first radial velocities for the system. They used the results deduced from the analysis of radial velocities for the solution of the V -passband light curve. Since CCD observations enabled to subtract the contamination of the neighboring field star, they derived the more accurate absolute parameters for the individual components. After performing a series of light curve solution, they estimated an effective temperature of 117 500 K for the primary component and fixing this value for the final light curve solution they suggested a temperature of ~ 7300 K for the secondary component.

In a later study, Bell et al. (1994, hereafter BPH94) confirmed that the primary eclipse is total with a duration of ~ 14 min. They also obtained the spectroscopic observations of the system during primary minima and extracted the spectrum of secondary component. Applying Kurucz (1979) atmosphere models to the spectra they determined a secondary component's temperature of 6250 ± 250 K. Adopting this value for the secondary component temperature they revised the preliminary light curve analysis. Solutions indicated a primary component with a temperature of $87\,000 \pm 13\,000$ K for a limb-brightened (Bond & Grauer 1987) secondary component.

V477 Lyr, the eclipsing binary nucleus of the PN Abell 46, has a similar light curve to UU Sge, except that the eclipse is partial (Bond 1980). The system shows a very large reflection effect which creates sine-like distortion on the light curve with an amplitude of ~ 0.6 mag.

Although its eclipsing nature was reported more than two decades ago, a few light curve solutions of V477 Lyr have been published. Ritter (1986) presented a solution based on a light curve analysis by L.W. Twigg (communicated by Bond, 1981). This solution yielded an orbital inclination of $80^\circ.8$ with a mass ratio of $M_1/M_2 \sim 2.3$, radii of $R_1 \sim 0.09 R_\odot$ and $R_2 \sim 0.20 R_\odot$. The temperature of the primary component was taken as $T_1 \sim 105\,000$ K. Considering a typical mass value for the central star of a planetary nebula (Schönberner 1981), Ritter assumed a mass for the primary component as $M_1 \sim 0.6 M_\odot$, which yields a secondary component mass of $M_2 \sim 0.25 M_\odot$. Also, Bond & Grauer (1987) cited the results of a preliminary analysis by J. Kaluzny and a temperature of $60\,000$ K for the primary component was suggested.

The most recent and detailed solutions of both light and radial velocity curves were made by Pollacco & Bell (1994, hereafter PB94). They obtained the medium-resolution spectra and a new V -passband CCD photometry of the system and presented the first complete light curve of V477 Lyr. They performed a series of light curve solution for the effective temperature of primary component ranging from $30\,000$ to $130\,000$ K. Using the mass ratio ($M_2/M_1 = 0.29$) obtained from the radial velocities and adopting the limb-brightening for the secondary component, they achieved a best fit to the observations for the effective temperature of $60\,000$ K. Thus, the V -passband light curve solution yielded an inclination of $\sim 80^\circ.5$, and a temperature of ~ 5300 K for the secondary component.

In this paper, we present new two-colour CCD photometry of UU Sge and V477 Lyr. The aim is to constrain the effective temperatures of the components through simultaneous

solutions of the two-colour light curves. We will estimate the common envelope efficiencies, α_{CE} , and discuss reasons for the oversized secondaries that we find.

2 OBSERVATIONS AND DATA REDUCTION

2.1 UU Sge

We have obtained the Johnson V - and R -passband photometry of the system at the National Observatory of the Scientific and Technological Research Council of Turkey (TÜBİTAK-TUG) in Antalya. TUG is established on the mountain of Bakırlitepe (in the West Taurus Mountain ranges), at a height of about 2500 m. Observations were carried out using a 1.5-m Cassegrain telescope (RTT 150) equipped with 2048×2048 back illuminated CCD camera which has an area of $9'.1 \times 9'.1$ and a scale of $0.26''/\text{pixel}$. We observed the system on the nights between July 4 and 7 and on the nights of 2002 August 6, 7 and 11. Since the orbital period of UU Sge is relatively short (~ 11.5 hours) and the system has a very deep primary minimum (~ 4.5 mag), exposure times were limited between 60 and 120 s. The light curves were phased by using the ephemeris given by Kiss et al. (2000),

$$\text{HJD (Min.I)} = 2451766.5285 + 0.46506921 E \text{ d.}$$

UU Sge has a nearby optical companion $\sim 2''.8$ to the east of the binary nucleus. The visual magnitude of the companion is $V=15.87$ mag. (Ciardullo et al. 1999). The visual magnitudes of the binary system itself at maxima and in mid-eclipse are $V_{\text{max}}=14.67$ mag and $V_{\text{min}}=19.24$ mag (PB93), respectively.

The data were reduced using the appropriate tasks in IRAF¹. First a master bias was created and subtracted from the images. Then we used an averaged master flat-field, which we created using the flat frames taken both during the evening and morning twilight, and normalized all the images. Since UU Sge is located within a crowded region, images were also trimmed right after bias and flat-field corrections. After first reduction steps were completed, IRAF/DAOPHOT package was used to apply the crowded field photometry. Thereby, we were able to make profile fitting and PSF (Point Spread Function) modeling of the stars in the target field and subtract the light contamination of the nearby optical companion. We measured the PSF magnitudes of UU Sge and five nearby stars to apply differential photometry. These

¹ IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.




Figure 1. *V*- and *R*-passband light curves of UU Sge (filled and open circles, respectively).

five stars were chosen as comparison star candidates which were previously determined by PB93 (C1, C2, C3, C5 and C6: Fig.1, therein). After investigating all comparison candidates for a possible variability, we decided to use C3 as the comparison, and we determined the differential magnitudes for UU Sge as plotted in Fig.1.

We have determined the depth of the total eclipse to be 4.450 ± 0.075 and 3.764 ± 0.074 mag for *V*- and *R*-passband, respectively. The depth for the *V*-passband is somewhat larger than that of BPH94. The time difference between the fourth and first-contact points was determined as ~ 1.26 hour, and the duration of the total eclipse at the bottom of the primary minimum was estimated as ~ 13.4 min. Not only the short time length but also the integration time increase at the bottom of the total eclipse, with a relatively long uploading time of the camera, brought about to obtain only a few data points in conjunction. Moreover, the minimum brightness of UU Sge is close to the magnitude limit (in case of photometry) for a 1.5-m telescope, which led to have scattered data points at the primary conjunction with errors around as much as 0.06 mag.

2.2 V477 Lyr

Two-passband (Johnson *V* and *R*) CCD photometric observations of V477 Lyr were carried out using the same setup as described for UU Sge. We observed the system during the nights from 2002 August 6 to 10. V477 Lyr has a partial eclipse at the primary minimum and it is




Figure 2. *V*- and *R*-passband light curves of V477 (filled and open circles, respectively).

not as deep as that of UU Sge. We set the exposure time as 60 s during the overall phases. The ephemeris used to phase the data was taken from Kiss et al. (2000),

$$\text{HJD (Min.I)} = 2451766.4038 + 0.47172909E \text{ d.}$$

After bias subtraction and flat-fielding, the data reduction was performed applying the aperture photometry package of IRAF/APPHOT. The magnitudes of the stars including V477 Lyr and other five comparison candidates, which were previously found to be constant by PB94 (C1, C3, C4, C5 and C6: Fig.1, therein), were measured. The errors of the magnitude measurements at primary minima did not exceed 0.018 and 0.010 mag in *V*- and *R*-passband, respectively. The differential magnitudes were computed with respect to the comparison star C6. The light curves of the system obtained in *V*- and *R*-passband are shown in Fig.2.

3 REANALYSIS OF THE RADIAL VELOCITIES

The first radial velocities of UU Sge and V477 Lyr were previously obtained and published by PB93 and PB94, respectively. By using both absorption and emission lines (which move in anti-phase with the absorption lines and thought to originate on or very close to the surface of the secondary component) they measured the radial velocities of the systems. They used one of the spectra of the related object (UU Sge or V477 Lyr) as a template to calculate the cross-correlations. Therefore, as mentioned by the authors, only the estimates

Table 1. Results of the radial velocity analysis.

Targets	K_1 (<i>km/s</i>)	K_2 (<i>km/s</i>)	q	$asini$ ($R_\odot$)
UU Sge	83.8 ± 6.7	182.8 ± 7.4	0.458 ± 0.034	2.45 ± 0.09
V477 Lyr	51.9 ± 6.4	182.1 ± 7.1	0.285 ± 0.026	2.18 ± 0.09

of the velocity semi-amplitudes are meaningful while the systematic velocities of the systems remain unknown. That is why we were not able to make simultaneous light and velocity curve solutions. However, we reanalysed the radial velocities by using the corrected radial velocity data given by PB93 and PB94 in order to increase the precision of the parameters and used them as inputs during the simultaneous solution of the two-colour light curves. The revised results of the radial velocity curve solutions are given in Table 1. The parameters K_1 and K_2 stand for the velocity semi-amplitudes of the primary and secondary components, q is the mass ratio (M_2/M_1) and a is the projected semimajor axis of the related system's orbit.

4 SIMULTANEOUS LIGHT CURVE SOLUTIONS

4.1 UU Sge

We have performed a simultaneous solution of the V - and R -passband light curves by employing the updated (2003) version of Wilson-Devinney (Wilson & Devinney 1971, Wilson 1979, 1990, 2003) code (hereafter W-D). This program is written for the analysis of light and velocity curves of binary stars which have either circular or eccentric orbits. The program has different ‘modes’ in terms of Roche lobe filling conditions of the components. Examination of the light curve of UU Sge revealed a detached system and a circular orbit as expected for such a close binary. Therefore, we used ‘Mode 2’ which is developed for detached binaries. Since the reflection effect is highly strong as seen in the light curves, a detailed treatment to the reflection effect was applied during the solutions.

The close binary systems, such as PNNi or post-CE (post-common envelope) binaries (subdwarf or white dwarf (WD) primary + low mass secondary) have large effective temperature differences between the components and show extreme reflection effects in their light curves due to irradiation around the substellar points of the cool secondaries. The modeling of the light curves of such systems is quite difficult and requires special treatment. As dis-

cussed by Hilditch et al. (1996), the temperature on the irradiated hemisphere of the cool secondary star is highly enhanced, therefore, contrary to the expected values ($A_2=0.5$, as generally assumed for cool stars) an albedo around or greater than one ($A_2 \geq 1$) is needed for securing a good fit to the detailed shape of the reflection effect (see also Wood et al. 1993). Previous studies also showed that the negative values of the limb-darkening coefficients are needed (limb-brightening, e.g. PB94, BPH94) due to same reason of obtaining more precise modeling of the light curves. In fact, by adopting negative limb-darkening coefficients for the secondary components of UU Sge and V477 Lyr, BPH94 and PB94 achieved more reasonable light curve solutions for albedo values greater than one.

We have performed a number of solutions by taking into account the previous approaches to the light curve modeling of the similar systems. The mass ratio ‘ q ’ obtained from the radial velocity analysis, the gravity brightening exponents of the components ($g_1=1.0$ and $g_2=0.32$), and the bolometric albedo of the primary component ($A_1=1$) were held fixed in the simultaneous solutions. Our first attempt to solve the light curves by using the square-root limb-darkening coefficients did not yield a rational solution, thence we decided to use linear limb-darkening law. The wavelength-specific linear limb-darkening coefficients of the hot companion were adopted as 0.10 and 0.07 for $x_1(V)$ and $x_1(R)$, respectively. These values were assumed using the tables given by Díaz-Cordovés et al. (1995) and Claret et al. (1995). Linear bolometric limb-darkening coefficients for both components were adopted from Van Hamme (1993, note that the limb-darkening coefficients given in these tables are calculated for up to an effective Kurucz atmosphere models (Kurucz 1993) were used for both stars during the solutions. The adjusted parameters were the orbital inclination (i), the mean effective temperature of the secondary, (T_2 —not including the effects that may influence the shape of the light curve such as reflection or spots), surface potentials of the components (Ω_1, Ω_2), bandpass luminosity of the primary component (L_1), bolometric albedo of the secondary component (A_2), and the wavelength-specific linear limb-darkening coefficients of the secondary component ($x_2(V)$, $x_2(R)$: the input values were taken from Claret (1998)). The main criterion to secure a reasonable solution was to achieve a physically realistic value for the albedo of the secondary component ($0.5 \leq A_2 \leq 1.0$). This criterion allowed us to set a lower limit for the temperature of the primary component, T_1 , and solution trials were started with a temperature of $T_1=73\,000$ K. So as to find the best solution, trials were made between the temperatures of 73 000 and 85 000 K. The temperature of the secondary component was first adopted as 3500 K considering the expected value of such a low mass star. Iterations were carried out until

Table 2. Parameters obtained from the simultaneous light curve analysis of UU Sge and V477 Lyr.

Parameter	UU Sge	V477 Lyr
i ($^{\circ}$)	87.12 ± 0.19	80.33 ± 0.06
Ω_1	7.497 ± 0.040	13.123 ± 0.099
Ω_2	3.309 ± 0.009	2.738 ± 0.006
T_1 (K)	78 000	49 500
T_2 (K)	6136 ± 80	3874 ± 120
$L_1/(L_1 + L_2)_V$	0.9837 ± 0.0035	0.9940 ± 0.0029
$L_1/(L_1 + L_2)_R$	0.9673 ± 0.0026	0.9811 ± 0.0031
A_1	1.0	1.0
A_2	0.927 ± 0.037	0.998 ± 0.036
g_1	1.0	1.0
g_2	0.32	0.32
$\langle r_1 \rangle$	0.1423 ± 0.0013	0.0779 ± 0.0007
$\langle r_2 \rangle$	0.2277 ± 0.0010	0.2091 ± 0.0003
$x_1(\text{bol})$	0.616	0.616
$x_2(\text{bol})$	0.493	0.384
$x_1(V)$	0.100	0.192
$x_1(R)$	0.070	0.162
$x_2(V)$	0.478 ± 0.078	0.210 ± 0.051
$x_2(R)$	-0.148 ± 0.074	-0.121 ± 0.053
rss	0.007	0.023

the convergence was reached at the smallest value for the residual sum of squares (rss). Although the solution trials have yielded parameters that are capable of modeling the light curves almost equal in appearance, we have achieved the best fit around the secondary minimum for a primary temperature of 78 000 K and a secondary temperature of 6136 ± 80 K, where the rss value was at a minimum among other temperatures adopted in the solutions. Examination of $T_1 - rss$ variation yielded an approximate error of 3000 K for the hot component's temperature. The uncertainty in the temperature of the secondary component, 80 K, is the formal 1σ error from the solution. The corrected error can be estimated as 240 K based on the uncertainty of 3000 K in the effective temperature of the primary. The results of the best solution are given in Table 2 (adjusted parameters have standard errors provided by the W-D code). The light curves and the computed fits are plotted in Fig.3. Some of the astrophysical parameters for the components were calculated and are provided in Table 3.

The mass of the secondary component seems considerably low compared to the mass of a main-sequence star with the same temperature. If we assume that the mass of the secondary component has not changed dramatically by the end of the CE interaction and it is just a low-mass main-sequence star, then the temperature seems significantly higher than that expected from an M type dwarf. The possible reasons which could explain this contradiction are discussed in § 6.

Since the simultaneous two-colour light curve solution gives us an opportunity to constrain the temperatures of the components, we believe that the temperatures derived in this




Figure 3. *V*- and *R*-passband light curves of *UU Sge* with computed fits.

paper are more accurate than the temperatures reported in previous studies. For example, Walton et al. (1993) suggested a primary component temperature of 50 000 K by comparing the merged ultraviolet spectrum with Kurucz (1991) LTE model atmospheres. They also obtained the spectrum of the secondary component during the primary eclipse (with some contamination from the sdO star) and assuming that the secondary is a main sequence star they classified it as G7 with an uncertainty of three spectral subclasses. BPH94 also took some spectra of the cool component during the primary minima. They used *ATLAS6* computer code (Kurucz 1979) and generated several atmosphere models to find the best fit to the spectrum of the secondary. Under the assumption that the secondary is a main sequence star they determined a temperature of 6250 ± 250 K. They used this value to reanalyse the light curve of *UU Sge* obtained by PB93 and gave two different results for the primary component temperature as $57\,000 \pm 8000$ and $87\,000 \pm 13\,000$ K, in terms of limb-darkening and brightening approximation, respectively.

The limb-darkening coefficients of the secondary component obtained from the solution significantly differ from those given by Claret (1998). If we compare our values with published ones, we see that the *V*-passband linear limb-darkening coefficient corresponds to a temperature of $\sim 10\,000$ K. The *R*-passband linear limb-darkening coefficient has a negative value and has no reference in the published tables. The only explanation we could make for

such limb-darkening coefficients would be the reflection effect emanated from the overheated face of the secondary, which gives rise to the “limb-brightening effect”.

There are several studies concerning the limb-darkening coefficients of illuminated atmospheres (e.g. Peraiah 1983, Claret 2004, Alencar & Vaz 1999). According to Alencar & Vaz (1999), as the illumination flux increases, the temperature distribution in the atmosphere of the secondary component becomes more homogeneous, in other words, center-to-limb brightness variation or center-to-limb-darkening over the stellar disk is removed. The observed increase in the intensity of the brightness of the secondary star from its center to its limb may be explained in terms of horizontal energy transport (Bond & Grauer 1987). On the other hand, the limb-darkening/brightening coefficients for the systems that have strongly irradiated atmospheres still need further investigation.

4.2 V477 Lyr

We have analysed the V - and R -passband light curves of V477 Lyr using the same code as described in the previous section. A series of simultaneous solution trials for a large temperature range of the primary component were performed. In order to obtain the best and realistic solution, we took into account the same criterion for the albedo of the secondary component, as we did for UU Sge. This criterion enable us to adopt a starting point for the temperature of hot component as 45 000 K. The fixed and adjusted parameters were the same as described for UU Sge. Similarly, we first attempted to apply the square root limb-darkening law during the solutions, as suggested by several authors for the stars have temperatures higher than 8000 K (e.g. Claret 1998), but we could not reach a rational solution and decided to apply the linear limb-darkening law. The input linear limb-darkening coefficients for the primary and secondary components were taken from the tables published by Díaz-Cordovés et al. (1995), Claret et al. (1995) and Claret (1998). Solution trials were made for the temperatures between 45 000 and 80 000 K. The best solution was achieved for the temperature of 49 500 K, where the rss value was at a minimum. And, the error in the temperature of the hot component was estimated as 4500 K from the examination of $T_1 - rss$ variation. The observed and computed light curves are given in Fig.4 and the parameters are listed in Table 2.

The linear limb-darkening coefficients derived from the solution exhibit a noteworthy discrepancy from those published ones that mentioned above. The cool component acts like




Figure 4. *V*- and *R*-passband light curves of V477 Lyr with computed fits.

a very hot star due to the reflection effect caused by the subdwarf primary. The heating on the hemisphere of the secondary star that faces to hot primary is such vigorous that the irradiation affects the limb-darkening over the stellar surface. Considering the grazing nature of the secondary eclipse, Bond & Grauer (1987) made an interpretation on the effectiveness of the reflection effect on V477 Lyr and suggested that the horizontal energy transport in the atmosphere of the secondary component might be the reason for the brighter limb.

To determine the temperature of the subdwarf companion, Kaler (1983) computed the Zanstra temperatures using the H and He II nebular lines and constrained the primary component's temperature in between 32 000 and 60 000 K. In a most recent study, PB94 presented a new Zanstra analysis based on the H β and He II nebular lines and estimated the temperatures as 48 000 and 82 000 K, respectively. They also employed the R-technique of Schönberner & Drilling (1984) and derived a temperature of $63\,000 \pm 12\,000$ K for the primary star and considering the temperature range found from several methods they decided to adopt a temperature of $60\,000 \pm 10\,000$ K in the light curve analysis. However, the simultaneous two-colour photometry gives us an opportunity to derive more accurate and approximate values for the temperatures of the binary components, which we have found as 49 500 K (fixed) for the primary and 3874 ± 120 K for the secondary star. The uncertainty of 120 K is the formal 1σ error from the solution. The corrected error can be estimated as 350 K based on the uncertainty of 4500 K in the effective temperature of the primary. The

Table 3. Astrophysical data for UU Sge and V477 Lyr.

Parameter	UU Sge	V477 Lyr
M_1 ($M_\odot$)	0.628 ± 0.053	0.508 ± 0.046
M_2 ($M_\odot$)	0.288 ± 0.031	0.145 ± 0.021
R_1 ($R_\odot$)	0.349 ± 0.013	0.172 ± 0.007
R_2 ($R_\odot$)	0.558 ± 0.021	0.462 ± 0.019
T_1 (K)	$78\,000 \pm 3000$	$49\,500 \pm 4500$
T_2 (K)	6136 ± 240	3874 ± 350
$\log g_1$ (<i>cgs</i>)	5.151 ± 0.050	5.672 ± 0.053
$\log g_2$ (<i>cgs</i>)	4.404 ± 0.058	4.269 ± 0.073

computed astrophysical parameters are given in Table 3. In fact, the mass of the secondary component corresponds to an M type star (as in the case of UU Sge) and a lower temperature is required for a main-sequence star at the same mass. One of the reasons why we could not reach such an expected temperature is the limb-brightening phenomenon, which takes place on the overheated hemisphere of the cool components.

5 THE REFLECTION EFFECT AND THE ENHANCED EFFECTIVE TEMPERATURES OF THE SECONDARY COMPONENTS

The hot subdwarf components of UU Sge and V477 Lyr, which are also responsible for the excitation of the nebulae, overheat the hemispheres facing to the substellar points of the cool components and the reflection effect reveals itself on the light curves' maxima as a sine-like distortion with large amplitudes (Figs. 1 and 2). The irradiation on the secondary components significantly increases the temperature around the substellar points. To estimate the reflection rates we applied Fourier analysis to the out-of-eclipse variation of the light curves and the following truncated Fourier series was used to compute the Fourier coefficients,

$$l = A_0 + A_1 \cos\theta + A_2 \cos 2\theta + B_1 \sin\theta + B_2 \sin 2\theta. \quad (1)$$

In this equation, l is the normalized intensity, θ is the phase in degrees, A_0 is the normalization level for the intensity, A_1 and A_2 are the contribution to the reflection effect. B_1 and B_2 are the coefficients represent the other perturbations on the light curve. The normalized Fourier coefficients of UU Sge and V477 Lyr for both passbands are listed in Table 4. The B coefficients have significantly smaller values than other coefficients, which indicate that other perturbations have no remarkable influence on the light curves other than reflection effect.

We have computed the amplitudes of the sine-like distortions caused by the reflection effect by using the Fourier coefficients of related systems. The amplitudes for UU Sge were

Table 4. Normalized Fourier coefficients ($l=1$ at $\theta=90^\circ$).

Targets	passbands	A_0	A_1	A_2	B_1	B_2
UU Sge	V	1.0265	-0.1907	0.0312	0.0062	-0.0002
		± 15	± 22	± 22	± 15	± 17
	R	1.0251	-0.2147	0.0356	0.0110	0.0016
		± 13	± 19	± 18	± 12	± 14
V477 Lyr	V	1.0477	-0.2912	0.0540	0.0053	0.0027
		± 22	± 33	± 25	± 30	± 29
	R	1.0630	-0.3338	0.0617	-0.0011	0.0015
		± 14	± 20	± 17	± 19	± 18

found to be 0.396 mag and 0.446 mag in V - and R -passband, respectively. The amplitude found in V -passband light curve seems in very good agreement with that of derived by PB93. In the case of V477 Lyr, the amplitudes in V - and R -passband were estimated to be 0.588 mag and 0.665 mag, in fair agreement with that of obtained by PB94.

We have tried to estimate the maximum temperature values, T_{2h} , around the substellar points of the secondary components by employing the following equation;

$$T_{2h}^4 = \frac{1}{R_2^2 \phi^2} [10^{-0.4\Delta m} (R_1^2 T_1^4 + R_2^2 T_2^4) - R_1^2 T_1^4]. \quad (2)$$

Where R_1 , R_2 are the radii and T_1 , T_2 are the effective temperatures of the primary and the secondary components, respectively; Δm is the amplitude of the brightness variation from minimum ($\theta=0^\circ$, mid-primary) to maximum ($\theta=180^\circ$, mid-secondary) of the sine-like distortion, and ϕ is the approximate central angle viewed from the centre of the secondary, which corresponds to the irradiated area over the surface of the secondary star (see Exter et al. 2005). We have determined the V - and R -passband substellar-point temperatures of the cool components of UU Sge to be around 30 680 K and 31 800 K, and of V477 Lyr to be around 16 860 and 17 550 K, respectively. These estimates agree well with those of computed by the equation described in Exter et al. (2005).

6 OVERSIZED RADII OF THE SECONDARY COMPONENTS AND EVOLUTIONARY STATUS OF THE SYSTEMS

Radii of the secondary components of UU Sge and V477 Lyr have been found to be larger than those expected for main-sequence stars at the same masses. On the other hand, our targets are not the only examples for such systems that contain oversized secondaries. In this section, we discuss the possible reasons on the oversized nature of the late-type secondary components of our targets along with some other selected PNNi and post-CE systems (listed in Table 5). Although the mass of the secondary component of V471 Tau is almost three times

the masses of other secondaries studied here, we have especially included this well-studied system for comparison. In a study made by O’Brien et al. (2001), two possible reasons for the expanded radius of the K-type secondary of V471 Tau were accentuated: (1) because of the possible CE phase, which V471 Tau might have recently emerged from, the secondary star may be still out of thermal equilibrium or (2) the starspots covering the surface of the star prevent the convective energy from being transported, which as a result increase the radius of the star.

The CE evolution and its effect on the low-mass main-sequence secondary components are the phenomena that have been poorly understood both theoretically and observationally. If a close binary system, consisting of a WD primary and a red dwarf (RD) secondary, have once undergone a CE phase, the mass and the radius of the secondary component might have been changed due to mass transfer during the formation of the CE. Although there remain questions on whether the mass of the secondary component alters dramatically or does not change in a substantial manner, there are observational evidences which show the significant changes in the radius of the component, such as the systems given in this paper.

There are several theoretical studies that have focused upon the response of the secondary component to the accretion process during the CE evolution. Prialnik & Livio (1985) made some evolutionary calculations for a $0.2 M_{\odot}$ fully convective star which accretes matter at several rates and investigated the outcome of the accretion. They found that in the models with accretion rates around 10^{-4} – $10^{-5} M_{\odot} \text{ yr}^{-1}$, the incoming material accumulates faster than the contraction rate, and consequently the outer radius increases. In their models the final value of the accumulated matter reaches to $2.5 \times 10^{-3} M_{\odot}$, which basically indicates that the mass of the fully convective star remains almost unchanged. They also concluded that the results should be valid for all fully convective stars ($\lesssim 0.3 M_{\odot}$) as well as for the stars with deep convective envelopes ($0.3 M_{\odot} \lesssim M \lesssim 0.5 M_{\odot}$). In a similar study carried out by Sarna & Ziolkowski (1988), the effect of rapid mass accretion onto low-mass main-sequence stars ($0.3 M_{\odot}$ – $0.5 M_{\odot}$) was investigated. They, too, came to a similar conclusion and indicated that the stars with deep convective envelopes experiencing rapid mass accretion significantly increase their radii. Another study published by Hjellming & Taam (1991) also probed the response of a $1.25 M_{\odot}$ main-sequence star to the conditions within the CE. At a range of relatively high accretion rates (10^{-5} to $10^{-1} M_{\odot} \text{ yr}^{-1}$), they found that the radius of the secondary component can expand ten times larger than its main-sequence radius upon mass accretion of less than $0.1 M_{\odot}$. After filling its Roche lobe, the secondary loses back

almost all the mass it has accreted and only a small amount of mass ($\lesssim 0.01 M_{\odot}$) remains accumulated after emergence from the envelope. As a result, the surface layers of the secondary depart from thermal equilibrium and enter in a thermally relaxation stage with a radius larger than its initial main-sequence radius. On the other hand, the results of some three-dimensional hydrodynamical simulations (e.g. Sandquist et al. 1998) reveal that the evolution of the CE phase takes place in short dynamic time scales and the secondary star which moves rapidly (supersonic) in the CE leaves this stage almost without mass accretion. However, the cases we will discuss in the following paragraphs involve the possible small amount of mass accretion by the cool companion.

The second reason which has been considered as a process to increase the radii of the low-mass secondary components is the effectiveness of the magnetic activity. There are several studies on low-mass main-sequence stars, which compare the physical properties of magnetically active and inactive stars. Mullan & MacDonald (2001) showed that, among the stars at the same masses, active stars are cooler and have larger radii than inactive stars. In a most recent study Morales et al. (2008) concluded that all active low-mass stars (including low-mass eclipsing binaries) are cooler than inactive stars of similar luminosity and hence have larger radii.

Recently, De Marco et al. (2008) have discussed the importance of the irradiation effect in binary PNNi and suggested that the heating which takes effect on the ‘day side’ of the secondary stars might cause a puffing up and expand the surface layers of the secondaries.

By taking into consideration two of the reasons summarized above, we have carried out some calculations to study the evolutionary status of some selected post-CE systems and interpreted which reason may be responsible for the cool components to have oversized radii: incomplete thermal relaxation or presence of magnetically active regions.

KV Vel : The binary nucleus of the PN DS 1. The sdO star was discovered by Drilling (1983) and double-lined spectroscopic variations with a period of 8.571 hour were found by Drilling (1985). Later, the renewed results for the system parameters were published by Hilditch et al. (1996).

BE UMa : It is an eclipsing binary nucleus. Although the light variability of the system was first noted by Kurochkin (1964) and an eclipse was first observed by Ando et al. (1982), the nebula surrounds the binary nucleus was discovered by Liebert et al. (1995). The system parameters listed in Table 5 and 6 have been adopted from Ferguson et al. (1999).

HS 1136+6646 : Discovered as a spectroscopic binary by Heber et al. (1996) and more

recently Sing et al. (2004) have indicated that the double-lined spectroscopic system is a newly formed post-CE binary.

PG 1026+002 : The system, which consists of a WD primary and a RD secondary, was discovered during the Palomar Green Survey (Green et al. 1986). Saffer et al. (1993) reported that the system is a single-lined spectroscopic binary and the secondary component's radius is compatible with its mass. However, by using the *R*-passband light curve of the system Bruch & Diaz (1999) found the radius of the secondary as being larger than expected from the theoretical models.

RR Cae : An eclipsing WD–M dwarf binary system with a period of 7.3 hour. The eclipses were first observed by Krzeminski (1984). Bruch (1999) analysed the radial velocity measurements together with the photometric data of Bruch & Diaz (1998) and found that the radius of the M dwarf is larger as compared with a main-sequence star at the same mass. However, in a more recent study, Maxted et al. (2007) presented new photometry and double-lined spectroscopy of the system and concluded that, contrary to Bruch (1999), the radius of the secondary is consistent with its mass. By taking into account the discrepancy between these two results we have performed two separate calculations.

RE 1016-053 : Post-CE binary consists of a WD primary and a RD secondary. Two different values for the secondary star's radius were given by Wood et al. (1999) and Vennes et al. (1999) depending on two different spectral analysis and the radius of the secondary component was found as slightly oversized. Both radii have been considered here and the results are listed in Table 5.

V471 Tau : Is a well-known post-CE binary system (WD primary + RD secondary) and a member of the Hyades star cluster. The system has a period of 12.5 hour and being observed since the discovery of its eclipses by Nelson & Young (1970). İbanoğlu et al. (1994) showed that the RD secondary is an active star. Possible reasons for the expanded radius of the secondary component were discussed by O'Brien et al. (2001) and they attributed the expansion to the magnetic activity which originates from the starspots partially covering the surface of the star.

We have used the models (age of 1 Gyr, $[M/H]=0.0$) provided by Chabrier & Baraffe (1997, hereafter CB97) to find the “*expected*” radii, temperatures and luminosities of the secondary components. The models have been fitted by sixth-degree polynomials and these equations have been used to calculate the “*expected*” radii ($R_{2,teo}$) given in Table 5 (along with “*observed*” ones). The mass–radius relationship of CB97 is plotted in Figure 5 and




Figure 5. Mass–radius relationships for the low mass main–sequence stars adopted from different studies. The fits are taken from: solid–line: CB97; dotted–line: Caillault & Patterson (1990); dashed–line: Lacy (1977). Radii of the secondary components are plotted as: UU Sge: filled diamond; V477 Lyr: open diamond; KV Vel: cross; BE UMa: asterisk; HS 1136+6646: open square; PG 1026+002: plus; RR Cae (Bruch 1999): filled circle; RR Cae (Maxted et al. 2007): open circle; RE 1016-053 (Wood et al. 1999): filled triangle; RE 1016-053 (Vennes et al. 1999): open triangle.




Figure 6. Mass–effective temperature relationship for the low mass main–sequence stars (CB97). The symbols are the same as in Figure 5.

comparison is made with the models of Lacy (1977) and Caillault & Patterson (1990). The ratio of the “*observed*” and “*expected*” radii are also given in Table 5. As it is seen from these ratios, most of the secondary components, including UU Sge and V477 Lyr, have relatively larger radii (2 or higher) than the radii of their main–sequence equivalents.




Figure 7. Mass–luminosity relationship for the low mass main–sequence stars. The symbols are the same as in Figure 5.

The locations of the secondary companions are illustrated in the mass–temperature (M – T_e) and mass–luminosity (M – L) planes in Figs. 6 and 7. Since the effective temperatures could not be determined for the secondary stars of HS 1136+6646 and RE 1016-053, the suggested spectral types taken from the referred papers are given in Table 5. In order to plot these stars in the M – T_e and M – L planes we adopted the effective temperatures from the tables given by Leggett (1992). The temperatures of 3800 K for K7 classification of HS 1136+6646, 3300 K and 3100 K for M3 and M5 classifications of RE 1016-053 were used in our calculations.

In order to have an idea on the thermal relaxation time–scale (t_{th}) of the envelope (accumulated matter) which was formed during the mass transfer from the primary to the secondary, we have used t_{th} equation given by Prialnik & Livio (1985):

$$t_{\text{th}} \simeq GMM_{\text{env}}\Delta R/R^2L. \quad (3)$$

Where M and R is the secondary’s mass and “*expected*” radius, M_{env} is the mass of the envelope due to mass accretion, ΔR is the width of the envelope and L is the bolometric luminosity of the secondary star.

Following the theoretical studies mentioned in earlier paragraphs we have assumed three M_{env} values which may have been accumulated by the secondary star: 0.05, 0.01 and 0.005 $M_{\odot}$. Then we have calculated the t_{th} for each assumption. The results are given in Table 5 as $t_{\text{th},0.05}$, $t_{\text{th},0.01}$ and $t_{\text{th},0.005}$.

Table 5. The observed and theoretical properties of the secondary components of the selected post-CE systems. The description of the properties are made in this section.

Object	M_2 ($M_\odot$)	$R_{2,\text{obs}}$ ($R_\odot$)	T_2 (K)	$T_{2,\text{teo}}$ (K)	$R_{2,\text{teo}}$ ($R_\odot$)	$R_{2,\text{obs}}/R_{2,\text{teo}}$	$t_{\text{th},0.05}$ (yr)	$t_{\text{th},0.01}$ (yr)	$t_{\text{th},0.005}$ (yr)	Ref.
UU Sge	0.288	0.558	6138	3420	0.280	1.989	1.64×10^8	3.27×10^7	1.64×10^7	this work
V477 Lyr	0.145	0.463	3874	3130	0.170	2.727	9.32×10^8	1.86×10^8	9.32×10^7	this work
KV Vel	0.230	0.402	3400	3340	0.238	1.688	1.62×10^8	3.24×10^7	1.62×10^7	1
BE UMA	0.360	0.720	5800	3480	0.336	2.142	1.28×10^8	2.56×10^7	1.28×10^7	2
HS 1136+6646	0.330	0.790	K7	3460	0.312	2.530	2.01×10^8	4.03×10^7	2.01×10^7	3
PG1026+002	0.220	0.400	3170	3330	0.231	1.733	1.85×10^8	3.71×10^7	1.85×10^7	4,5
RR Cae	0.095	0.189	2500	2730	0.118	1.598	1.07×10^9	2.15×10^8	1.07×10^8	6
RR Cae	0.182	0.210	3100	3250	0.202	1.042	1.45×10^7	2.90×10^6	1.45×10^6	7
RE 1016-053	0.285	0.357	M3	3420	0.278	1.283	4.76×10^7	9.53×10^6	4.76×10^6	8
RE 1016-053	0.150	0.265	M5	3150	0.174	1.521	2.62×10^8	5.24×10^7	2.62×10^7	9
V471 Tau	0.930	0.960	5040	5370	0.832	1.153	5.15×10^5	1.03×10^5	5.15×10^4	10

References: The references are given for the observed properties of the secondary components. (1) Hilditch et al. (1996), (2) Ferguson et al. (1999), (3) Sing et al. (2004), (4) Bruch & Diaz (1999), (5) Saffer et al. (1993), (6) Bruch (1999), (7) Maxted et al. (2007), (8) Wood et al. (1999), (9) Vennes et al. (1999), (10) O’Brien et al. (2001).

Table 6. The properties of the primary components of the selected post-CE systems.

Object	M_1 ($M_\odot$)	T_1 (K)	$\log g_1$ (cgs)	Age (yr)	Ref.
UU Sge	0.628	78 000	5.151	3.0×10^4	1
V477 Lyr	0.508	49 500	5.672	3.0×10^4	1
KV Vel	0.63	77 000	5.85	3.0×10^4	1
BE UMA	0.70	105 000	6.5	3.0×10^4	1
HS 1136+6646	0.63	70 000	7.75	9.0×10^5	2
PG1026+002	0.65	17 600	8.1	1.4×10^8	2
RR Cae	0.467	7005	7.7	1.9×10^9	3
RR Cae	0.44	7540	7.7	1.9×10^9	3
RE 1016-053	0.57	55 800	7.81	1.8×10^6	2
RE 1016-053	0.59	55 000	7.84	1.8×10^6	2
V471 Tau	0.84	34 500	8.31	1.2×10^7	2

References: The references given here are for the models which have been used to compute the “age” of the systems (for the other properties given in this table please see the references given in Table 5). (1) Schönberner (1983), (2) AB98, (3) BA98.

Since it is very important to know the critical accretion rates, we have also calculated the Eddington luminosities of the secondary stars. We have found the critical accretion rates to be of the order of a few $10^{-4} M_\odot \text{ yr}^{-1}$. Therefore, it is acceptable to consider a mass accretion onto secondary components as rapid as a few $10^{-5} M_\odot \text{ yr}^{-1}$, which is an adequate accretion rate leads to increase in the radii of the cool components as previously discussed by Prialnik & Livio (1985).

For the binary nuclei, UU Sge, V477 Lyr, KV Vel and BE UMa, the presence of the PN warrants us in estimating the ages (the time since the emergence from the CE) of the systems as a few 10^4 yr (30 000 – 40 000 yr, Schönberner 1983, Soker 2006). So, we have simply adopted the ages as $3 \times 10^4 \text{ yr}$ for all the nuclei (Table 6). However, we have used the evolutionary tracks computed by Blöcker (1995), and showed the possible evolutionary states of the sdO componets of UU Sge and V477 Lyr in Fig. 8. Although these models




Figure 8. The evolutionary status of UU Sge and V477 Lyr. The models are taken from Blöcker (1995). Open and filled circles represent the primary component of UU Sge and V477 Lyr, respectively.

were computed for PNe with single nuclei, the loci of the sdO components seem in fairly well agreement (within the uncertainties given for the masses of sdO components) with the models have remnant masses of 0.625 and 0.524 $M_{\odot}$, which evolved from the initial ZAMS masses of 3 and 1 $M_{\odot}$, respectively. In these models, the remnants have CO cores and the nuclear energy is mainly provided by the He shell burning. Besides all these approaches, we are aware of the fact that our systems are not PNe with “single” nuclei, and binarity may have influenced the evolution of the components through the interaction between the components and formation of the CE (more details on the evolution of PNe and CE can be found in the referred papers), as much as the different mass-loss rates the systems might have encountered.

For the other systems studied here we have used the white dwarf evolutionary models from Althaus & Benvenuto (1998, hereafter AB98) and Benvenuto & Althaus (1998, hereafter BA98) to calculate the approximate cooling ages, t_{cool} . In AB98, the evolutionary models were computed for the carbon–oxygen WD masses within the range of 0.5–1.2 $M_{\odot}$ and surface hydrogen layer masses ranging between 10^{-13} and $10^{-4} M_{\text{H}}/M$, where M is the mass of the modeled WD and M_{H} is the mass of the H envelope. BA98 developed the models for the evolution of helium WDs with masses ranging from 0.15 to 0.5 $M_{\odot}$, and having outer H envelope masses within the range of $10^{-8} \leq M_{\text{H}}/M \leq 4 \times 10^{-3}$. And both WD models were improved for the metallicities of $Z=0.0$ and $Z=0.001$.

For the primary masses higher and lower than $0.5 M_{\odot}$, we have used the tables given by AB98 and BA98, respectively. Various M_{H}/M and both Z values have been taken into account and the models have been examined in $T_{\text{e}}\text{--}\log g$ plane. The most applicable models have been selected and interpolated for the related primary masses in order to estimate the cooling ages of the WDs. We have found that the models for V471 Tau and RR Cae having the surface hydrogen layer mass of $10^{-4} M_{\text{H}}/M$ give the best agreement in $T_{\text{e}}\text{--}\log g$ plane for the metallicities of $Z=0.0$ and $Z=0.001$, respectively. However, other systems seem consistent with the models have surface hydrogen layer mass of $10^{-6} M_{\text{H}}/M$ and metallicity of $Z=0.001$. Resultant t_{cool} values are listed in Table 6.

Since our goal is to search for the more plausible reason on the oversized nature of the secondaries, we have compared t_{th} with the ages of the systems as summarized below:

UU Sge, *V477 Lyr*, *KV Vel* and *BE UMa* : The thermal relaxation time-scales calculated for each of the envelope masses are relatively longer than the ages of the systems, which indicate that the surface of the secondary components have not yet reached the thermal equilibrium. The discrepancy between the “*observed*” and the “*expected*” radii of the secondaries can be seen in Fig. 5. The difference of the effective temperatures from the theoretical ones is also notable (Fig. 6).

HS 1136+6646 : The comparison of t_{th} and t_{cool} of the system shows that surface layers of the system is still in a thermally relaxation stage which causes the expansion in the radius of the secondary. Also the adopted effective temperature which corresponds to a K7 star is located above the theoretical fit and appears to be in agreement with this approach.

PG 1026+002 : When the computed t_{th} values are compared with t_{cool} of the system, depending on the envelope mass, the secondary component seems to be either at the end of its thermally relaxation stage or just passed it. However, the low effective temperature of the secondary component may be taken as support for the idea that the reason for the expanded radius is the presence of the possible magnetically active regions on the secondary star.

RR Cae : For the parameters given by Bruch (1999), the thermal relaxation time-scales for the secondary component of RR Cae are shorter than the cooling age of the system. The inconsistency both in radius and effective temperature with the theoretical expectations indicates the presence of the possible stellar activity. On the other hand, the possibility of the higher amount of matter (at least $0.1 M_{\odot}$) being accreted by the secondary leaves a question mark that the cool component might be still out of thermal equilibrium. Maxted et al. (2007)

found the radius consistent with the theoretical expectations as seen in Fig. 5. But there seems to be a disagreement between effective temperature and its expected counterpart, which also supports the assumption of the presence of active regions. Because the mass of the secondary component is twice as much of the mass found by Bruch (1999), t_{th} values are relatively well below the cooling age of the system.

RE 1016-053 : There is an interesting result for the secondary star of RE 1016-053. The radius of the cool component (taken from both Wood et al. 1999 and Vennes et al. 1999) has larger values than expected according to the mass–radius relationship of CB97 and the comparison of the thermal relaxation time–scales with the cooling age of the system shows that the secondary star still may have thermal instability on its surface layers. But the adopted effective temperatures seem to be a little lower than expected, which could tell us the possibility of having active regions on the surface of the secondary if we would have lower t_{th} values than t_{cool} . However, our calculations indicate a secondary star which is far from being thermally relaxed that makes us to expect higher effective temperatures. Since the effective temperatures have been determined from the spectral types estimated by the authors (Wood et al. 1999 and Vennes et al. 1999) and they suggest these spectral types relying on the masses of the secondary star, we believe this is due to probable underestimation of either spectral types or the masses.

V471 Tau : Our findings on this well-known post-CE system are listed in Table 5 and 6, but since the mass of the secondary component is higher than the rest of the secondaries studied here, it is not plotted in Figs. 5, 6 and 7. It is obvious that the secondary component passed its thermally relaxation stage relatively long ago. Nonetheless, its radius seems a little larger and the effective temperature seems a little lower. These results are in fairly well agreement with those presented by O’Brien et al. (2001).

7 THE EFFICIENCY PARAMETER : “ α_{CE} ”

The CE evolution is one of the most crucial part in the evolution of binary systems. The presence of close binaries, for example, consisting of a WD primary and a RD secondary, indicates that the wide precursor system must have undergone a significant amount of angular momentum loss which causes the orbital shrinkage.

At some stage of the evolution, more massive primary component evolves to AGB (or RGB, depending on the initial configuration of the system) and fills its Roche lobe which

leads to mass transfer towards the secondary component. If the mass transfer is unstable the secondary star can not accommodate the accreted matter at its surface and both components are engulfed by the primary's envelope which is later on being called as CE. Since the CE does not corotate with the orbiting binary core, the frictional forces arise due to velocity difference. This frictional interaction produces gravitational drag that causes components to spiral-in together and reduces the binary separation. The change in orbital separation results in the change in orbital energy and a fraction of the difference in orbital energy is used to eject the envelope from the system by depositing it into CE. This fraction is designated as the “*efficiency*” parameter and denoted by “ α_{CE} ” (e.g. Tutukov & Yungelson 1979, Iben & Livio 1993, and references therein)

$$\alpha_{\text{CE}} = \frac{\Delta E_{\text{bind}}}{\Delta E_{\text{orb}}}, \quad (4)$$

where ΔE_{bind} is the difference between gravitational and thermal energy of the ejected envelope at the beginning of Roche lobe overflow (RLOF) and ΔE_{orb} is the orbital energy difference between the beginning and at the end of the spiralling-in process of the binary system.

One of the main goals of this paper is to make a contribution to the studies which search for the limits to constrain the value of “ α_{CE} ” parameter. We have mainly followed the approximations given by, e.g., Webbink (2008), Han et al. (1995), Dewi & Tauris (2000), De Kool (1990) in order to estimate the “*efficiency*” parameters for UU Sge and V477 Lyr. We assume that the secondary components are main-sequence stars and have not been affected by the CE interaction. For the estimation of the binding energy of the envelope (ΔE_{bind}) we have used the following equation

$$\Delta E_{\text{bind}} = -\frac{GM_1 M_{\text{env}}}{\lambda R_L}, \quad (5)$$

where M_1 is the mass of AGB star (also assumed as the initial mass of the primary component), $M_{\text{env}} = M_1 - M_{\text{core}}$, is the envelope mass of AGB star, M_{core} is the current mass of the primary, R_L is the Roche lobe radius of the primary (at the onset of mass transfer) and λ is a dimensionless parameter which depends on the detailed structure of the envelope and the stellar density distribution. According to Webbink (2007), λ is a function of $m_e \equiv M_{\text{env}}/M_1 = 1 - (M_{\text{core}}/M_1)$, and is well-approximated by

$$\lambda^{-1} \approx 3.000 - 3.816m_e + 1.041m_e^2 + 0.067m_e^3 + 0.136m_e^4. \quad (6)$$

The total change in orbital energy is

$$\Delta E_{\text{orb}} = \frac{GM_{\text{core}}M_2}{2a_f} - \frac{G(M_{\text{core}} + M_{\text{env}})M_2}{2a_0}, \quad (7)$$

where M_2 is the mass of secondary component, a_0 is the initial separation of the binary system, and a_f is the final orbital separation.

In order to calculate these energies one needs to estimate the initial parameters of the binary systems. As a first conjecture for the initial masses of the primaries we have made use of the evolutionary tracks of Blöcker (1995). However, by taking into consideration that the RLOF may occur at different stages of the binary evolution we have also used the initial/final mass relations given by Iben & Tutukov (1986) (also see Iben & Livio 1993). If the components experience a case-C RLOF (primary fills its Roche lobe during the He-shell burning) then the mass of the remnant is related to initial mass of the primary by $M_{\text{CO}} \approx 0.5 + 0.16(M_1 - 2.3)$ if $2.3 < M_1 < 4.2$ (the masses are in solar units) and by $M_{\text{CO}} \approx 0.8 + 0.0875(M_1 - 4.2)$ if $4.2 < M_1 < 7.6$. If the binary system experiences a late case-B RLOF (primary fills its Roche lobe shortly before the He ignition, He core is non-degenerate at this point), a degenerate CO or a “hybrid (C+O+He)” remnant develops with an initial/final mass relation of $M_{\text{CO}} \approx 0.167M_1 - 0.085$ if $2.3 < M_1 < 5$, and $M_{\text{CO}} \approx 0.4 + 0.07M_1$ for the range of $5 < M_1 < 10$.

Considering the different stages of RLOF mentioned above, let us first investigate the case-C RLOF in which the primary components of UU Sge and V477 Lyr are CO cores. Then the initial masses of the primary stars can be estimated as $\approx 3.1 M_{\odot}$ and $\approx 2.35 M_{\odot}$, respectively. If we assume that the systems have experienced a case-B RLOF and become degenerate CO or hybrid cores, then the initial masses of the primary components must have been around $4.2 M_{\odot}$ and $3.5 M_{\odot}$, respectively. By taking these masses as limits for the progenitors of the primaries we have performed a series of calculations to estimate the α_{CE} parameters of the systems (following a somewhat similar work to that of O’Brien et al. 2001).

We used the theoretical AGB core mass-luminosity approximation given by Iben & Tutukov (1984, 1985, 1993);

$$L_{\text{AGB}} = 6 \times 10^4 (M_{\text{CO}}/M_{\odot} - 0.5), \quad (8)$$

where L_{AGB} is the maximum luminosity for a post-AGB star during the plateau phase of its evolution and also the luminosity at the time of RLOF. Applying this approximation, L_{AGB} values for the current primary masses of UU Sge and V477 Lyr can be calculated as 7680 and $480 L_{\odot}$, respectively.

Table 7. Possible initial configurations of the systems and the efficiency parameter, α_{CE} .

Object	M_1 ($M_\odot$)	R_{AGB} ($R_\odot$)	q	a_0 ($R_\odot$)	P_i (<i>days</i>)	α_{CE}
UU Sge	3.0	297	10.42	511	738	0.48
	3.1	294	10.76	503	710	0.51
	4.2	266	14.58	438	501	0.99
V477 Lyr	1.0	60	6.90	110	125	0.71
	2.35	45	16.21	73	47	4.86
	3.5	40	24.14	62	30	12.82
V477 Lyr*	2.2	210	15.17	345	485	0.83

* The sdO component has been assumed as a degenerate “He” dwarf.

The AGB radii of the primaries have been estimated using the AGB mass–luminosity–radius relation proposed by Hurley et al. (2000);

$$R_{\text{AGB}} = 1.125M_1^{-0.33}(L_{\text{AGB}}^{0.4} + 0.383L_{\text{AGB}}^{0.76}), \quad (9)$$

where R_{AGB} is the AGB radius that the progenitor might have reached at the stage of RLOF. The R_{AGB} values calculated for the different initial masses are also given in Table 7.

Now we can estimate the initial separations, a_0 , of the systems with the following formula (Eggleton 1983)

$$r_L = \frac{0.49q^{2/3}}{0.6q^{2/3} + \ln(1 + q^{1/3})}, \quad (10)$$

where $r_L = R_L/a_0$ is the relative radius of the Roche lobe and q is the mass ratio being equal to M_1/M_2 . Adopting $R_L = R_{\text{AGB}}$ the initial separations have been computed for different mass ratios (see Table 7). Equations (5) and (6) have been used for the estimation of the binding and orbital energies, then the α_{CE} parameters have been determined as listed in Table 7. P_i , the initial orbital period of the system has been also computed using the Kepler’s third law and given in Table 7.

The somewhat problematic point here is that the mass of the sdO component of V477 Lyr is of the order of $0.5 M_\odot$ which is a critical and controversial value that lies in the transition region between the masses of CO and He remnants. This situation gives us a chance to consider the sdO component as a degenerate “He” dwarf which once have a progenitor mass less than $2.3 M_\odot$. If we assume that the primary component filled its Roche lobe right after it has developed a degenerate “He” core, then we can employ the core–mass–radius relation given by Iben & Tutukov (1986),

$$R = 10^{3.5} M_{\text{He}}^4. \quad (11)$$

Here, R is the radius of primary by the time of RLOF and M_{He} is the current mass of the

sdO star. By adopting this radius as R_L and assuming that the initial mass of the primary was around $2.2 M_\odot$ we have calculated the initial parameters and values of α_{CE} for the system as listed in Table 7.

8 SUMMARY AND DISCUSSION

In this paper, we report the results of the first simultaneous two-passband light curve analysis for the binary PNNi; UU Sge and V477 Lyr. We have also reanalysed the published radial velocity data (PB93 and PB94) of both systems in order to increase the sensitivity of the orbital parameters employed in the solution of the light curves. The linear limb-darkening coefficients were adjusted during the solutions of both systems. Not surprisingly, the resultant coefficients were found to be inconsistent with the theoretically expected ones, which was interpreted as the presence of the limb-brightening effect due to strong irradiation of the secondary components. In the case of UU Sge, the best solution was achieved at the primary temperature of 78 000 K by yielding a secondary component temperature of 6136 K. The temperature of the primary component falls into the temperature range of $87\,000 \pm 13\,000$ K derived by BPH94. Also the temperature of secondary component indicates a good agreement with that of 6250 K estimated by BPH94.

We have determined the temperatures of the hot and cool components of V477 Lyr as 49 500 K and 3874 K, respectively. The temperature of the secondary star shows a remarkable discrepancy in comparison with the temperature of 5300 K derived by PB94. We believe that having two-passband simultaneous light curve analysis with a different approach to the solutions yielded more realistic results in terms of the temperature of the components of both systems.

Nevertheless, it has to be stressed that the secondary component temperature of V477 Lyr, in contrary to that of UU Sge, seems much closer to the value of expected from a main-sequence star at the same mass. The discrepancy between the observed and theoretical temperatures may be explained in terms of the thermally relaxation stage that the secondary stars have been shown to be in. The more the mass is accumulated the longer the thermal-relaxation time scale is needed. During the computations, our main assumption was that the mass of the secondary stars has not been significantly affected by the CE interaction and the highest value for the mass of the accumulated matter was adopted as $0.05 M_\odot$. However, we have to bring into attention that the mass of the secondary components might

have undergone drastic changes due to accretion, which would also result in the increase of the luminosity (and the temperature) of the secondary components.

The temperature of the primary component of V477 Lyr indicates a good agreement with the $H\beta$ Zanstra temperature of 48 000 K while showing a disagreement with the He II Zanstra temperature of 82 000 K, those of derived by PB94. 49 500 K also lies in the range of H and He II Zanstra temperatures of 32 000 K – 60 000 K determined by Kaler (1983). PB94 mentions the possible He II stratification in the nebula, which leads up to the variety in the He II Zanstra temperatures of the sdO component. The mass loss via thermal pulses may explain the likely He stratification and enhanced He abundance (PB94) in the nebula. In this case, the Zanstra temperature determined from He II nebular lines could be misleading, which reveals that the temperature determined by using the H nebular lines may carry more valuable information on the temperature of the sdO component.

The comparison of the thermal time scales and the post-CE ages of the systems (Table 5) indicates that the secondary components of the binary nuclei of the PNe are still out of thermal equilibrium along with two other post-CE systems: HS 1136+6646 and RE 1016-053. The rest of the systems that have been reported as having oversized secondary components were found to be thermally relaxed relatively long ago. The oversized nature and low temperatures of these systems were interpreted as the presence of the possible magnetically active regions which cause the increase in the size of a low-mass main sequence star while reducing its effective temperature. Detailed spectroscopic studies of these systems are needed.

Finally, we have implemented several calculations to estimate the common envelope efficiency parameter, α_{CE} , of the systems. We have determined various initial system parameters using possible evolutionary scenarios. The possible efficiency parameters for UU Sge were found in the range of $0.3 < \alpha_{\text{CE}} < 1$, which is close to the range of previously suggested by theoretical studies (e.g. Han et al. 1995). Nevertheless, the parameter values that we have found for V477 Lyr, under the assumption of having CO or C+O+He remnant as the primary component, indicate a deviation from the theoretical findings. An $\alpha_{\text{CE}} > 1$ may be explained as the contribution from other sources (e.g. deposition of recombination energy into CE ejection, nuclear burning on the surface of the secondary star, see Iben & Livio (1993) for more detail) to the CE ejection beside the efficiency of orbital energy. On the other hand, it seems impossible to know which mechanism could have been such effective on the efficiency parameter. Therefore, we have found the courage of making a new assump-

tion about the primary component of V477 Lyr. By taking into account its current mass of $0.508 \pm 0.046 M_{\odot}$, we assumed that the primary component is a degenerate He dwarf with an initial mass of $< 2.3 M_{\odot}$. By using the mass–radius relation given by Iben & Tutukov (1986), we calculated the radius of the primary at the time of RLOF (independently from its luminosity, see equation (11)). These approaches yielded an efficiency parameter of 0.83 which indicates that about 83 percent of the orbital energy were used effectively during the ejection of the CE.

MA would like to thank the Scientific and Technological Research Council of Turkey (TÜBİTAK) and her advisor, Cİ, for their support during her doctoral studies. The authors are grateful to Howard E. Bond for his help and useful suggestions on this manuscript. We also wish to thank TÜBİTAK National Observatory of Turkey (TUG) for the allocation of observing time. And we thank the referee very much for the comments and suggestions that have helped us to improve the manuscript. This work has been supported by the Ege University Research Foundation under the project number of 2002/FEN/004.

REFERENCES

- Abell G. O., 1966, *ApJ*, 144, 259
- Afşar M., Bond H. E., 2005, *MmSAI*, 76, 608
- Alencar S. H. P., Vaz L. P. R., 1999, *A&AS*, 135, 555
- Althaus L. G., Benvenuto O. G., 1998, *MNRAS*, 296, 206
- Ando H., Okazaki A., Nishimura S., 1982, *PASJ*, 34, 141
- Bell S. A., Pollacco D. L., Hilditch R. W., 1994, *MNRAS*, 270, 449
- Benvenuto O. G., Althaus L. G., 1998, *MNRAS*, 293, 177
- Blöcker T., 1995, *A&A*, 299, 755
- Bond H. E., 1976, *PASP*, 88, 192
- Bond H. E., Liller W., Mannery E. J., 1978, *ApJ*, 223, 252
- Bond H. E., 1980, *IAU Circ.*, 3480
- Bond H. E., Grauer A. D., 1987, in *IAU Colloq. 95, Second Conference on Faint Blue Stars* (Dordrecht: Kluwer), 221
- Bond H. E., Livio M., 1990, *ApJ*, 355, 568
- Bond H. E., 2000, in *Kastner J. H., Soker N., Rappaport S., eds, ASP Conf. Ser. Vol. 199*,

- Asymmetrical Planetary Nebulae II: From Origins to Microstructures. Astron. Soc. Pac., San Francisco, p. 115
- Bruch A., Diaz M. P., 1998, AJ, 116, 908
- Bruch A. 1999, AJ, 117, 3031
- Bruch A., Diaz M. P., 1999, A&A, 351, 573
- Budding E., Kopal Z., 1980, Ap&SS, 73, 83
- Caillault A. P., Patterson J., 1990, AJ, 100, 825
- Chabrier G., Baraffe I., 1997, A&A, 327, 1039
- Ciardullo R., Bond H. E., Sipior M. S., Fullton L. K., Zhang C.-Y., Schaefer K. G., 1999, AJ, 118, 488
- Claret A., 1998, A&A, 335, 647
- Claret A., Díaz-Cordovés J., Giménez A., 1995, A&AS, 114, 247
- Claret A., 2004, A&A, 422, 665
- De Kool M., 1990, ApJ, 358, 189
- De Marco O., Bond H. E., Harmer D., Fleming A. J., 2004, ApJ, 602, L93
- De Marco O., Hillwig, T. C., Smith A. J., 2008, AJ, 136, 323
- Dewi J. D. M., Tauris T. M., 2000, A&A, 360, 1043
- Díaz-Cordovés J., Claret A., Giménez A., 1995, A&AS, 110, 329
- Drilling J. S., 1983, ApJ, 270, L13
- Drilling J. S., 1985, ApJ, 294, 107
- Eggleton P. P., 1983, ApJ, 268, 368
- Exter K. M., Pollacco D. L., Maxted P. F. L., Napiwotzki R., Bell S. A., 2005, MNRAS, 359, 315
- Ferguson D. H., Liebert J., Haas S., Napiwotzki R., James T. A., 1999, ApJ, 518, 866
- Green R. F., Schmidt M., Liebert J., 1986, ApJS, 61, 305
- Han Z., Podsiadlowski P., Eggleton P. P., 1995, MNRAS, 272, 800
- Heber U., Dreizler S., Hagen H. J., 1996, A&A, 311, L17
- Hilditch R. W., Harries T. J., Hill G., 1996, MNRAS, 279, 1380
- Hjellming M. S., Taam, R. E., 1991, ApJ, 370, 709
- Hoffleit, D., 1932, Harvard Bull., No. 887
- Hurley J. R., Pols O. R., Tout C. A., 2000, MNRAS, 315, 543
- İbanoğlu C., Keskin V., Akan M. C., Evren S., Tunca Z., 1994, A&A, 281, 811
- Iben I. Jr., Tutukov A. V., 1984, ApJS, 54, 335

- Iben I. Jr., Tutukov A. V., 1985, *ApJS*, 58, 661
- Iben I. Jr., Tutukov A. V., 1986, *ApJ*, 311, 753
- Iben I. Jr., Livio M., 1993, *PASP*, 105, 1373
- Iben I. Jr., Tutukov A. V., 1993, *ApJ*, 418, 343
- Kaler J. B., 1983, *ApJ*, 271, 188
- Kiss L. L., Kasza J., Borza S., 2000, *IBVS*, 4962
- Krzeminski W., 1984, *IAU Circ.* 4014
- Kurochkin N. E., 1964, *Peremnye Zvezdy* (Academy Sci. U.S.S.R., Var. Star Bull), 15, 77
- Kurucz, R. L., 1979, *ApJS*, 40, 1
- Kurucz R. L., 1991, in Davis Philips A. C., Upgren A. R., James K. A., eds, *Proc. Workshop on Precision Photometry: Astrophysics of the Galaxy*. Schenectady, NY, p. 27
- Kurucz R. L., 1993, in Milone E. F., ed., *Light Curve Modelling of Eclipsing Binary Stars*. Springer-Verlag, New York, p. 93
- Lacy C. H., 1977, *ApJS*, 34, 479
- Leggett S. K., 1992, *ApJS*, 82, 351
- Liebert J., Tweedy R., Napiwotzki R., Fulbright M. S., 1995, *ApJ*, 441, 424
- Maxted P. F. L., ODonoghue D., Morales-Rueda L., Napiwotzki R., Smalley B., 2007, *MNRAS*, 376, 919
- Morales J. C., Ribas I., Jordi C., 2008, *A&A*, 478, 507
- Mullan D. J., MacDonald J., 2001, *ApJ*, 559, 353
- Nelson B., Young A., 1970, *PASP*, 82, 699
- O'Brien M. S., Bond H. E., Sion E. M., 2001, *ApJ*, 563, 971
- Paczynski B., 1976, in *IAU Symp. 73, Structure and Evolution of Close Binary Systems*, ed. P. Eggleton, S. Mitton, J. Whelan (Dordrecht: Reidel), 75
- Peraiah A., 1983, *JApA*, 4, 11
- Pollacco D. L., Bell S. A., 1993, *MNRAS*, 262, 377
- Pollacco D. L., Bell S. A., 1994, *MNRAS*, 267, 452
- Prialnik D., Livio M., 1985, *MNRAS*, 216, 37
- Ritter H., 1986, *A&A*, 169, 139
- Saffer R. A., Wade R. A., Liebert J., Green R. F., Sion E. M., Bechtold J., Foss D., Kidder K., 1993, *AJ*, 105, 1945
- Sandquist E. L., Taam R. E., Chen X., Bodenheimer P., Burkert A., 1998, *ApJ*, 500, 909
- Sarna M. J., Ziółkowski J., 1988, *AcA*, 38, 89

- Schönberner D., 1981, A&A, 103, 119
- Schönberner D., 1983, ApJ, 272, 708
- Schönberner D., Drilling J. S., 1984, ApJ, 278, 702
- Sing D. K., Holberg J. B., Burleigh M. R., Good S. A., Barstow M. A., Oswalt T. D., Howell S. B., Brinkworth C. S. et al., 2004, AJ, 127, 2936
- Soker N., 2006, ApJ, 645L, 57
- Tutukov A. V., Yungelson L. R., 1979, in Mass loss and evolution of O-type stars, ed. C. de Loore, P. S. Conti (Reidel, Dordrecht) 401
- Van Buren D., Charles P. A., Mason K. O., 1980, ApJ, 242L, 105
- Van Hamme W., 1993, AJ, 106, 2096
- Vennes S., Thorstensen J. R., Polonski E. F., 1999, ApJ, 523, 386
- Walton N. A., Walsh J. R., Pottasch S. R., 1993, A&A, 275, 256
- Webbink R. F., 2008, Short-Period Binary Stars: Observations, Analyses, and Results. Series: Astrophysics and Space Science Library, XVIII, Vol. 352, p.233. Edited by E.F. Milone, D.A. Leahy, and D.W. Hobill
- Wilson R. E., 1979, ApJ, 234, 1054
- Wilson R. E., 1990, ApJ, 356, 613
- Wilson R. E., Devinney E. J., 1971, ApJ, 166, 605
- Wilson R. E., Van Hamme W., 2003, Documentation of Eclipsing Binary Computer Model. Dept. Astron., Univ. Florida, Gainesville, FL, USA
- Wood J. H., Zhang E-H., Robinson E. L., 1993, MNRAS, 261, 103
- Wood J. H., Harmer S., Lockley J. J., 1999, MNRAS, 304, 335

This paper has been typeset from a $\text{\TeX}$ / $\text{\LaTeX}$ file prepared by the author.